

Incomplete completions of local rings

September 2000

Let A be a local ring (commutative with 1) with maximal ideal \mathfrak{m} . The projection maps $A \rightarrow A/\mathfrak{m}^n$ for $n = 1, 2, \dots$, induce a local homomorphism of local rings

$$c_A : A \longrightarrow \widehat{A} = \varprojlim_n A/\mathfrak{m}^n.$$

We say that A is complete if c_A is an isomorphism. We denote the maximal ideal of \widehat{A} by $\widehat{\mathfrak{m}}$. Let M_i be the kernel of the surjective projection map $\widehat{A} \rightarrow A/\mathfrak{m}^i$.

Proposition 1 *The local ring \widehat{A} is complete if and only if $M_n = \widehat{\mathfrak{m}}^n$ for each $n \geq 1$.*

Proof. For each n the projection map $\widehat{A} \rightarrow A/\mathfrak{m}^n$ gives rise to a short exact sequence

$$0 \longrightarrow (M_n/\widehat{\mathfrak{m}}^n) \longrightarrow (\widehat{A}/\widehat{\mathfrak{m}}^n) \longrightarrow (A/\mathfrak{m}^n) \longrightarrow 0.$$

If we let n run this becomes a short exact sequence of projective systems. The projection map $\widehat{A} \rightarrow A/\mathfrak{m}^{n+1}$ sends $\widehat{\mathfrak{m}}^n$ onto $\mathfrak{m}^n/\mathfrak{m}^{n+1}$, so $\widehat{\mathfrak{m}}^n + M_{n+1} = M_n$. This implies that the system on the left has surjective transition maps. By Mittag-Leffler we get a short exact sequence of projective limits:

$$0 \longrightarrow \varprojlim_n M_n/\widehat{\mathfrak{m}}^n \longrightarrow \widehat{A} \xrightarrow{g_A} \widehat{A} \longrightarrow 0.$$

It follows that g_A is an isomorphism if and only if all $M_n/\widehat{\mathfrak{m}}^n$ are zero. But we have $g_A c_{\widehat{A}} = \text{id}_{\widehat{A}}$, so g_A is an isomorphism if and only if $c_{\widehat{A}}$ is an isomorphism. \square

Proposition 2 *For any local ring A for which \widehat{A} is not complete, the ring $\widehat{\widehat{A}}$ is not complete either.*

Proof. For each $n \geq 1$ we consider the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \varprojlim_n M_n/\widehat{\mathfrak{m}}^n & \longrightarrow & \widehat{\widehat{A}} & \xrightarrow{g_{\widehat{A}}} & \widehat{A} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & M_n/\widehat{\mathfrak{m}}^n & \longrightarrow & \widehat{A}/\widehat{\mathfrak{m}}^n & \longrightarrow & A/\mathfrak{m}^n & \longrightarrow & 0 \end{array}$$

in which the rows are exact and the vertical maps are surjective. By the snake lemma the kernel K_n of the middle vertical map maps surjectively to M_n . Thus, K_n/K_1^n surjects to M_n/M_1^n , and by the previous criterion, \widehat{A} is not complete. \square

The next result is exercise 12 in Bourbaki, Commutative Algebra, Ch. II sec. 2 (p. 235/236).

Proposition 3 *There is a local ring A so that \widehat{A} is not complete.*

Proof. Let K be a field. Let A_d be the localisation of $K[X_1, \dots, X_d]$ at the maximal ideal (X_1, \dots, X_d) , and let A be the union of all A_d . Let \mathfrak{m} be the maximal ideal of A and let \mathfrak{m}_d be the maximal ideal of A_d . Note that \widehat{A} is the power series ring $K[[X_1, X_2, \dots]]$ in which all elements have only finitely many terms of each total degree. In particular, for any monomial in the variables X_1, X_2, \dots we can consider the coefficient of an element of \widehat{A} at that monomial.

We will show that $M_2 \neq \widehat{\mathfrak{m}}^2$.

First note that for $a < b$ a polynomial map from affine a -space over K to affine b -space over K is not surjective on K -rational points (finite K : cardinality; infinite K : algebraic geometry).

We will construct a sufficient condition for an element of M_2 not being an n -term sum of products of two elements of $\widehat{\mathfrak{m}}$. For each d and $n \geq 2$ we look at the image of a polynomial map of affine spaces over K :

$$f_d : (\mathfrak{m}_d/\mathfrak{m}_d^n \times \mathfrak{m}_d/\mathfrak{m}_d^n)^n \longrightarrow \mathfrak{m}_d^n/\mathfrak{m}_d^{n+1},$$

that sends $(x_i, y_i)_{i=1}^n$ to the degree n part of $\sum_i x_i y_i$. For fixed n the K -dimension of the affine space on the right is a polynomial of degree n in d with a positive leading coefficient and on the left it is a polynomial of degree $n - 1$ in d . (In fact, on the left it is $2n \binom{d+n-1}{n-1} - 1$, and on the right it is $\binom{d+n-1}{n}$.) For given n this implies that for sufficiently large d there is a homogeneous polynomial s_n of degree n in d variables which is not in the image of f_d . By considering the map $\widehat{A} \rightarrow A_d/\mathfrak{m}_d^{n+1}$ that sends all variables X_i with $i > d$ to zero we see that any power series in M_2 which has s_n as its degree- n -part is not a sum of n products of pairs of elements of $\widehat{\mathfrak{m}}$. By summing the elements s_n over all $n \geq 2$ we find an element of M_2 that is not in $\widehat{\mathfrak{m}}^2$. \square

Bart de Smit
desmit@math.leidenuniv.nl

H. W. Lenstra, Jr.
hwl@math.berkeley.edu